

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.e-jds.com

Correspondence

Pathological mandibular fractures following idiopathic resorption of bilateral mandibular rami and coronoid processes

KEYWORDS

Coronoid process;
Idiopathic mandibular
resorption;
Mandibular ramus;
Pathological fracture

Pathological mandibular fractures are rare, accounting for fewer than 2% of all mandibular fractures.¹ Pathological fractures usually may follow surgical interventions such as third molar removal or implant placement, and are caused by osteomyelitis, osteonecrosis, cyst, or tumor.¹ Therefore, idiopathic mandibular fracture is rarer, especially the bilateral fractures. Here, we reported a rare case of pathological mandibular fractures following idiopathic resorption of bilateral mandibular rami and coronoid processes.

A 74-year-old man with open bite and bilateral mandibular pain and swelling was referred to our department. Panoramic radiograph showed bilateral mandibular fractures caused by resorption of the bilateral mandibular rami and coronoid processes (Fig. 1A). Panoramic radiograph taken 20 months before the initial visit revealed resorption of the bilateral mandibular angles (Fig. 1B). Three-dimensional computed tomography (3D-CT) showed open bite and bilateral mandibular fractures (Fig. 1C and D). Bimaxillary removable partial prostheses were fabricated to improve occlusion. Although several examinations were performed to identify the cause of mandibular resorption, systemic sclerosis with oral manifestations of malocclusion and mandibular resorption was ruled out. Panoramic radiograph taken 2 months after the initial visit revealed further resorption of the bilateral mandibular fracture sites (Fig. 1E). However, there was slight

resorption of the mandibular fracture sites in 3D-CT at 6 months after the first visit (Fig. 1F and G). Therefore, mandibular reconstructive surgery will be performed, when mandibular resorption does not occur after the follow-up of more than 1 year from the first visit.

Progressive osteolysis or resorption of the mandible is a rare entity. Mandibular resorption in systemic sclerosis predominantly occurs at areas of muscle attachment, such as the mandibular angle, coronoid process, and mandibular condyle.² The current theory behind the etiology of bone resorption is multifactorial, including ischemia secondary to pressure necrosis from fibrosis and atrophic musculature, as well as decreased vascularity as a result of vessel wall sclerosis.² Gorham's disease (GD) is an extremely rare bone disorder characterized by osteolytic bony resorption and can affect any bone in the body.³ It is of unknown etiology and thought to result from localized endothelial proliferation of lymphatic vessels that cause bony resorption.³ The diagnosis can only be made after excluding other common causes of bony resorption such as those caused by infection, inflammation, malignancy, and endocrine.³ In maxillofacial GD, the mandible is more commonly affected than the maxilla.⁴ Maxillofacial GD can affect any age group, and most cases are detected before 40 years of age, with a mean presenting age of 29.6 years.⁴ The mobility of teeth can be the initial presenting finding, and the jaw continues to resorb and eventually disappear completely.⁴ The

<https://doi.org/10.1016/j.jds.2023.09.021>

1991-7902/© 2023 Association for Dental Sciences of the Republic of China. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Figure 1 (A) Panoramic radiograph at the initial visit, (B) Panoramic radiograph at 20 months before the initial visit, (C, D) Three-dimensional computed tomography at the initial visit, (E) Panoramic radiograph at 2 months after the initial visit, (F, G) Three-dimensional computed tomography at 6 months after the initial visit.

differential diagnosis includes Paget disease of bone, Langerhans cell histiocytosis, osteomyelitis, primary intra-osseous malignancy, metastatic malignancy, hyperparathyroidism, aneurysmal bone cyst, and rheumatoid arthritis.⁴

In the present case of an elder, spontaneous resorption of bilateral mandibular angles and coronoid processes resulted in a pathological mandibular fracture. However, the cause of the mandibular resorption was not clear, because the patient refused a biopsy for the diagnosis. Once bilateral mandibular resorption has stopped, we will plan mandibular reconstruction with the biopsy.

Conflicts of interest

The authors have no conflicts of interest relevant to this article.

References

1. Boffano P, Rocca F, Gallesio C, Berrone S. Pathological mandibular fractures: a review of the literature of the last two decades. *Dent Traumatol* 2013;29:185–96.
2. Burchfield C, Vorrasi J. Maxillofacial implications of scleroderma and systemic sclerosis: a case report and literature review. *J Oral Maxillofac Surg* 2019;77:1203–8.
3. Kunjur J, Hirani M, Manisali M, Matthews NS. Total joint replacement for severe bilateral condylar resorption- Gorham-Stout syndrome. *J Craniofac Surg* 2015;26:e302–5.
4. Sansare K, Saalim M, Jogdand M, Karjodkar FR, Sreenivasan V. Radiographic extent of maxillofacial Gorham's disease and its impact on recurrence: a systematic review. *Oral Surg Oral Med Oral Pathol Oral Radiol* 2021;132:80–92.

Haruka Yoshii
Toshinori Iwai*
Satomi Sugiyama
Kenji Mitsudo

Department of Oral and Maxillofacial Surgery/
Orthodontics, Yokohama City University Hospital,
Yokohama, Kanagawa, Japan

*Corresponding author. Department of Oral and Maxillofacial Surgery/Orthodontics, Yokohama City University Hospital, 3-9 Fukuura, Kanazawa-ku, Yokohama, Kanagawa 236-0004, Japan.

E-mail address: iwai104@yokohama-cu.ac.jp (T. Iwai)

Received 18 September 2023

Final revision received 18 September 2023

Available online 28 September 2023