

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.e-jds.com

Original Article

TNF- α promotes expression of inflammatory factors by upregulating nicotinamide adenine dinucleotide phosphate oxidase-2 expression in human gingival fibroblasts

Min Xu^{a†}, Churen Zhang^{b†}, Ye Han^a, Jie Zhang^a,
Xiaochi Chang^a, Jianxia Hou^{a*}, Sha Li^{c**}

^a Department of Periodontology, National Center of Stomatology & National Clinical Research Center for Oral Diseases, National Engineering Laboratory for Digital and Material Technology of Stomatology, Peking University School and Hospital of Stomatology, Beijing, China

^b Department of Stomatology, The First Affiliated Hospital of Xiamen University, School of Medicine, Xiamen University, Xiamen, China

^c College of Pharmacy, Jinan University, Guangzhou, China

Received 22 April 2023; Final revision received 24 April 2023

Available online 7 May 2023

KEYWORDS

Inflammation;
Human gingival
fibroblasts (HGFs);
Nicotinamide adenine
dinucleotide
phosphate oxidase-
2 (NOX-2);
Tumor necrosis
factor- α (TNF- α)

Abstract *Background/purpose:* Periodontitis is a chronic infectious disease. The oxidative stress environment can cause or exacerbate the inflammation in periodontitis. Nicotinamide adenine dinucleotide phosphate oxidase (NOX) may be the most important source of reactive oxygen species (ROS) in periodontal tissues. The pathological mechanism of periodontitis may be related to the increased ROS caused by enhanced NOX activity. The purpose was to investigate the effect of tumor necrosis factor (TNF- α) on inflammatory cytokines and ROS, and the role of NOX-2 in human gingival fibroblasts (HGFs).

Materials and methods: HGFs were cultured and divided into the normal control group (NC group) and the inflammatory model group (TNF- α group) induced by 10 ng/ml TNF- α . Thereafter, NOX-2 siRNA was used to knock down NOX-2 gene expression. Quantitative real-time PCR was applied to detect *IL-6*, *MCP-1*, and *NOX-2* mRNA levels. The levels of *IL-6* and *MCP-1* protein were examined by ELISA. The level of NOX-2 was evaluated by Western blot. ROS expression was measured by the fluorescence microplate.

Results: The mRNA and protein expression levels of *IL-6*, *MCP-1*, and *NOX-2* were significantly increased, and the expression of ROS was significantly elevated in response to 10 ng/ml TNF- α .

* Corresponding author. Department of Periodontology, Peking University School and Hospital of Stomatology, No. 22, Zhongguancun South Avenue, Haidian District, Beijing, 100081, China.

** Corresponding author. College of Pharmacy, Jinan University, No. 601, West Huangpu Avenue, Guangzhou, 510632, Guangdong, China. E-mail addresses: jxhou@163.com (J. Hou), tlisha@jnu.edu.cn (S. Li).

† These two authors contributed equally to this work.

Compared with the si-NC group, the mRNA and protein expression levels of *IL-6* and *MCP-1* were significantly down-regulated and ROS expression was significantly decreased in the si-NOX2 group stimulated by 10 ng/ml TNF- α .

Conclusion: TNF- α promotes the expression of NOX-2 in human gingival fibroblasts and enhances the expression of inflammatory factors and ROS in human gingival fibroblasts through the upregulation of NOX-2 partly.

© 2023 Association for Dental Sciences of the Republic of China. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Periodontitis is a chronic infectious disease caused by plaque microorganisms. The interaction between the plaque microorganisms and the host determines the development and process of the disease, leading to the inflammatory destruction of periodontal tissue.¹ The main clinical manifestations of periodontitis include gingival redness, attachment loss, formation of the periodontal pocket, alveolar bone resorption, and tooth loosening and displacement. It is the leading cause of tooth loss in adults.² In recent years, increasing evidence has shown that the oxidative stress environment can cause or exacerbate the inflammatory response of periodontitis.³

Reactive oxygen species (ROS) are considered a “double-edged sword” that exerts its effects by fighting against microorganisms. Under physiological conditions, ROS resist and kill invading pathogenic microorganisms.⁴ ROS play an important role in cell signaling, gene regulation, and antimicrobial defense.⁵ However, excessive ROS production can lead to increased oxidant load with unchanged or reduced antioxidant capacity, resulting in a state of oxidative stress in the tissues. It can cause pathological changes that damage the periodontal supporting tissues of the host and eventually lead to tooth loss with the destruction of periodontal tissue structure.^{6,7} There has been a longstanding interest in the relationship between ROS and periodontitis.^{8,9}

Nicotinamide adenine dinucleotide phosphate oxidase (NOX) may be the most important source of ROS in periodontal tissues.¹⁰ NOX is the terminal component of the respiratory chain and plays a role in electron transfer, obtaining a single electron from the cytoplasmic nicotinamide adenine dinucleotide phosphate (NADPH), transferring it through the plasma membrane to external oxygen molecules, and leading to the production of ROS.¹¹ It has been suggested that the pathological mechanism of periodontitis may be related to the increased production of ROS caused by enhanced NOX activity. Therefore, it is crucial to inhibit NOX activity and adequately regulate the production of ROS in periodontitis.

Seven isoforms of NOX proteins exist in human organs, namely NOX 1–5, DUOX-1, and DUOX-2.¹² The NOX protein family is a multimeric complex (excluding NOX-5) that includes transmembrane subunits (such as NOX-2, p22^{phox}) and cytoplasmic subunits (such as p47^{phox}, NOXO1, p67^{phox}, NOXA1 p40^{phox}, Rac).¹⁰ In the resting state, the cytoplasmic

and transmembrane subunits are separated. In the activated state, the cytoplasmic subunits undergo phosphorylation and translocation to the transmembrane subunits, forming a polymeric complex that functions.¹³ NOX-2 is expressed in the gingival tissue of chronic periodontitis.¹⁴ For example, in NOX-2, p47^{phox}, p40^{phox}, and p67^{phox} exist as a complex in the cytoplasm in the resting state. In the activated state, the p47^{phox} subunit is phosphorylated and then this complex as a whole moves toward the plasma membrane and binds to Cytb558 on the membrane, forming a NOX polymer complex.¹⁵

Gingival fibroblasts (GFs) have the ability to proliferate and self-renew rapidly.¹⁶ The maintenance of normal physiological functions such as proliferation and differentiation of GFs is an important mechanism for self-healing after periodontal tissue injury.¹⁷ In this study, we investigated the effects of TNF- α on inflammatory cytokines and ROS in human gingival fibroblasts and the role of NOX-2 through *in vitro* experiments.

Materials and methods

Primary cell culture

Human gingival fibroblasts (HGFs) were extracted from healthy gingival tissues. Gingival tissues were obtained from healthy periodontal individuals requiring crown lengthening surgery in the Department of Periodontology, Peking University School and Hospital of Stomatology, Beijing, China. The Biomedical Ethics Committee of Peking University School and Hospital of Stomatology approved this study (IRB00001052-08010), and informed consent forms were signed prior to the surgery. Gingival tissues were immediately immersed in Dulbecco Modified Eagle Medium (DMEM; Gibco, Grand Island, NY, USA) supplemented with penicillin (200 U/ml)–streptomycin (200 U/ml) (Gibco). After removal of the epithelium, collected gingival specimens were minced into small pieces (0.5–1 mm³ in size), and then plated onto a cell culture dish in DMEM medium containing 10% Foetal Bovine Serum (FBS; Gibco) and penicillin (100 U/ml)–streptomycin (100 U/ml). The cells were incubated at 37 °C in a 5% CO₂, 95% humidified atmosphere. The following experiments were conducted with 3–6 passages of HGFs. HGFs were divided into two groups: the normal control group (NC group) and the inflammatory cell group (TNF- α group). TNF- α (PeproTech, Cranbury, NJ,

USA) was added to the TNF- α group when the cell density reached 80%–90% (final concentration of 10 ng/ml). Cells were continued to be cultured for 48 h, and samples were collected.

Small interfering RNA transfection

NOX-2-specific small interfering RNA (siRNA) and non-targeting control siRNA were synthesized by RiboBio (Guangzhou, China). 1×10^6 HGFs were seeded into 6-well plates (Corning Inc., Rochester, NY, USA) for 24 h, and the siRNAs were transfected into HGFs grown to 70% using Lipofectamine 3000 (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instructions. The sequence of NOX2 siRNA was 5'-CCGGAGGTCTTACTTTGAA-3'.

Quantitative real-time polymerase chain reaction

Cells were collected and Trizol reagent (Ambion, Austin, TX, USA) was added to extract total RNA from cells. Total RNA content was detected by spectrophotometer. RNA was reversely transcribed into cDNA using reverse transcription kit (TOYOBO, Osaka, Japan) and cDNA was used as the template for real-time fluorescence quantitative PCR detection (TOYOBO). Using GAPDH as internal reference, mRNA relative expression levels of IL-6, MCP-1, and NOX family (including NOX-1, NOX-2, NOX-3, NOX-4, and NOX-5) were calculated by the $2^{-\Delta\Delta C_t}$ method. The primers were synthesized by Sangon Biotech (Shanghai, China) as follows: 5'-CGACAGTCAGCCGCATCTT-3' (forward primer) and 5'-CCAA-TACGACCAAATCCGTTG-3' (reverse primer) for the human GAPDH gene, 5'-GTGAGGAACAAGCCAGAGC-3' (forward primer) and 5'-TACATTTGCCGAAGAGCC-3' (reverse primer) for the human IL-6 gene, 5'-CAGCCAGATGCAATCAATGCC-3' (forward primer) and 5'-TGGAATCCTGAACCCACTTCT-3' (reverse primer) for the human MCP-1 gene, 5'-GGAAT-TAGGCAAAGTGGGTTTT-3' (forward primer) and 5'-CAGTG GCCTTGTCAAAGTTTAA-3' (reverse primer) for the human NOX-1 gene, 5'-AAGATGCGTGGAAGTACCTAA-3' (forward primer) and 5'-CAGTGGCCTTGTCAAAGTTTAA-3' (reverse primer) for the human NOX-2 gene, 5'-AACATCACCTTCTG-TAGAGACC-3' (forward primer) and 5'-CTTCTTGTTG AAATCGCCAGAA-3' (reverse primer) for the human NOX-3 gene, 5'-TCACAGCCTCTACATATGCAAT-3' (forward primer) and 5'-CAGCAGCATGTAGAAGACAAAG-3' (reverse primer) for the human NOX-4 gene, 5'-TGAGCAGAAAGACACTATCTGG-3' (forward primer) and 5'-CTGATGCCTTGAAGGACTCATA-3' (reverse primer) for the human NOX-5 gene.

Western blot analysis

Cells were collected and added with RIPA lysate (Solarbio, Beijing, China) and PIC protease inhibitor (Solarbio). Ultrasonic cracking and centrifugal collection of supernatant. The BCA protein detection kit (Thermo Fisher Scientific, Waltham, MA, USA) measured the protein concentration of the sample. The samples were separated by gel electrophoresis with sodium dodecyl sulfate polyacrylamide (SDS-PAGE; Dakewe Biotech, Shenzhen, China), after which the

proteins were transferred to polyvinylidene fluoride (PVDF) membranes (Thermo Fisher Scientific). 5% skim milk (BD, Franklin Lakes, NJ, USA) was enclosed for 1 h and the membrane was hybridized with NOX-2 monoclonal antibody (Abcam, Cambridge, UK) or GAPDH monoclonal antibody (Cell Signaling Technology, Beverly, MA, USA). TBST was washed and incubated with horseradish peroxidase labeled secondary antibodies (Cell Signaling Technology). ECL reagent (Solarbio) was used for chemiluminescence. The protein band strength was quantitatively analyzed by Image J software, and the GAPDH protein strength was used as an internal reference for standardized analysis.

Enzyme-linked immunosorbent assay

Cell supernatant of cultured HGFs was collected, centrifuged, and stored at -80°C . Content of IL-6 and MCP-1 in supernatant was detected by Enzyme-linked immunosorbent assay (ELISA) according to the instructions of the human IL-6 and MCP-1 ELISA kit (Meimian, Jiangsu, China).

Reactive oxygen species ROS

ROS levels were detected using a chemiluminescent probe, DCFH-DA reagent (Beyotime Biotechnology, Shanghai, China). Cells were stimulated with 10 ng/ml TNF- α for 24 and 48 h. Then serum-free DMEM medium containing DCFH-DA was added (volume ratio 1:1000) and cells were incubated at 37°C for 20 min. After trypsin digestion, cells were collected by centrifugation after suspension, washed with PBS 2–3 times, and cell precipitates were collected for fluorescence detection. The collected cells were made into cell suspension with PBS. The excitation wavelength was 488 nm while the emission wavelength was 525 nm.

Statistical analysis

Excel 2019, SPSS 25.0, and Prism 8.0 statistical analysis software were used to analyze the experimental data. Each experiment was repeated more than 3 times. The unpaired T test was used to compare the difference between the experimental and control groups. $*P < 0.05$ was considered statistically significant.

Results

TNF- α promoted expression of IL-6 and MCP-1 in human gingival fibroblasts

We firstly constructed an inflammatory cell model using TNF- α of 10 ng/ml. RT-qPCR was used to detect mRNA expression level of IL-6 and MCP-1 and our results showed that the expression level of IL-6 and MCP-1 in TNF- α group was significantly increased (Fig. 1A and B). Concentrations of IL-6 and MCP-1 in supernatant were detected by ELISA and the results demonstrated that TNF- α significantly up-regulated expression of IL-6 and MCP-1 protein in the supernatant (Fig. 1C and D).




Figure 1 Gene and protein expression levels in HGFs. (A) The relative mRNA expression of IL-6 was detected by RT-qPCR. (B) The relative mRNA expression of MCP-1 was detected by RT-qPCR. (C) Concentration of IL-6 in supernatant was measured by ELISA. (D) Concentration of MCP-1 in supernatant was measured by ELISA. *: $P < 0.05$; **: $P < 0.01$.

Expression of ROS was upregulated in human gingival fibroblasts with TNF- α stimulation

In order to elucidate the association between TNF- α and oxidative stress, we determined ROS levels in HGFs with TNF- α stimulation. Our results found that intracellular ROS levels markedly increased after incubation with 10 ng/ml TNF- α in HGFs for 48 h (Fig. 2A and B).

TNF- α promoted the expression of NOX-2 in human gingival fibroblasts

To determine the role of NOX in HGFs, we examined the mRNA expression of the five NOX family members in response to 10 ng/ml TNF- α . Among these NOX family mRNA, NOX-2 expression was significantly increased (Fig. 3B) and NOX-4 expression was significantly decreased (Fig. 3D) in the TNF-

α group of HGFs, while incubation of HGF with TNF- α had no significant effect on the mRNA expression of NOX-1, NOX-3, and NOX-5 (Fig. 3A, C, and E). The results of Western blot analysis demonstrated that the protein expression of NOX-2 was also significantly elevated in TNF- α group compared with the NC group in HGFs (Fig. 3F and G).

NOX-2 gene silencing reduced the expression level of IL-6 and MCP-1

NOX-2 siRNA (si-NOX2) and negative control siRNA (si-NC) were transfected with HGFs, followed by stimulation with 10 ng/ml TNF- α for 48 h. The mRNA expression level of IL-6 and MCP-1 was detected by RT-qPCR, and the protein level of IL-6 and MCP-1 in the cell culture supernatant were measured by ELISA. After knockdown of NOX-2, the mRNA expression levels (Fig. 4A and B) and protein secretion




Figure 2 Production level of ROS in HGFs. (A) The intracellular ROS (green) with or without 10 ng/ml TNF- α stimulation for 48 h. Bar = 100 μ m. (B) The intracellular ROS levels that HGFs produced as measured by fluorescence microplate. **: $P < 0.01$. Abbreviations: ROS, reactive oxygen species.

levels (Fig. 4C and D) of IL-6 and MCP-1 were significantly down-regulated in response of TNF- α .

NOX-2 gene silencing decreases the expression of ROS

The results of fluorescence microplate revealed that NOX-2 downregulation significantly decreased the level of ROS production in the si-NOX2 group compared with the si-NC group (Fig. 5A and B).

Discussion

Periodontitis is a chronic infectious disease with progressive loss of periodontal tissues caused by plaque microorganisms, eventually leading to tooth loosening and abscission.² In 2010, periodontitis was the sixth most common disease worldwide, with a prevalence of approximately 10.8%, affecting 743 million people worldwide.¹⁸ TNF- α is an important pro-inflammatory cytokine that plays an important role in the attachment loss of periodontal tissues and the progression of periodontal diseases.¹⁹ 10 ng/ml TNF- α is often used to induce inflammatory cell models.^{16,20} Therefore, in this study, 10 ng/ml TNF- α was used to induce an inflammation model of human gingival fibroblasts.

IL-6 is an inflammatory cytokine involved in bone resorption.²¹ MCP-1 is a chemokine secreted by various cell types involved in a variety of biological functions such as recruitment of inflammatory cells, wound healing, inhibition of stem cells, and maintenance of effector immune responses.^{22,23} Changes in IL-6 and MCP-1 levels can reflect the severity of periodontal inflammation.²⁴ Therefore, this study investigated the expression levels of inflammatory

cytokines IL-6 and MCP-1 in inflammatory models of human gingival fibroblasts. The results showed that the expression levels of genes and proteins of IL-6 and MCP-1 were significantly increased after 10 ng/ml TNF- α stimulation in human gingival fibroblasts. This proves that TNF- α upregulates the expression levels of IL-6 and MCP-1 in human gingival fibroblasts.²⁵

In recent years, many studies have shown that the oxidative stress environment is an important factor in the occurrence and progression of a large variety of chronic inflammatory diseases, such as periodontitis,⁵ type 2 diabetes,²⁶ atherosclerosis,²⁷ rheumatoid arthritis,²⁸ and inflammatory lung disease.²⁹ A large number of case-control and longitudinal studies comparing the levels of different antioxidant markers in the saliva of healthy and periodontitis subjects have shown that the levels of oxidative stress biomarkers (such as 8-OHdG and MDA) in saliva of periodontitis patients are significantly increased compared with healthy individuals.^{30–32} After non-surgical treatment for periodontitis, the levels of antioxidants in saliva of periodontitis patients were significantly increased and were associated with the improvement of periodontal clinical index after treatment.³³ These results indicate a direct correlation between increased oxidative stress or ROS levels and the development of periodontitis or clinical improvement of periodontitis after treatment.^{30,33}

As considered a "double-edged sword", ROS play an important role in combating microorganisms under physiological conditions, while cause cytotoxicity to host cells when overactivated.³⁴ The relationship between ROS and periodontitis has long been a concern.^{8,9} In recent years, NOX has been considered to be the main source of ROS in periodontal tissues. In this study, the 10 ng/ml TNF- α induced inflammation cell model showed that TNF- α can promote the production of ROS.³⁵ Similarly, compared with




Figure 3 Gene and protein expression levels of the NOX family in HGFs. (A–E) The relative mRNA expression of NOX-1, NOX-2, NOX-3, NOX-4, NOX-5 was detected by RT-qPCR. (F) The protein expression of NOX-2 and GAPDH in HGFs. (G) The relative protein level of NOX-2 compared with GAPDH in HGFs. *: $P < 0.05$; **: $P < 0.01$; ns: not significant.

the control group, the expression levels of the NOX-2 gene and protein in the inflammatory cell model group were significantly increased. This proves that TNF- α can upregulate the expression level of NOX-2 in human gingival fibroblasts. Lin et al. reported that TNF- α significantly promoted the activation of NOX and the production of ROS (including $O_2^{\cdot-}$ and H_2O_2), using TNF- α to induce human alveolar epithelial cells.³⁶ This is consistent with our findings.

ROS in periodontal tissue is believed to be mainly produced by NOX.³⁵ At present, there is limited research on the impact of NOX-2 on inflammatory cytokines IL-6 and MCP-1. Some studies have shown that in the mouse model of periodontitis induced by *Aggregatibacter actinomycetemcomitans* (Aa), NOX-2 gene knockout mice have

significantly reduced alveolar bone volume, increased inflammatory cell infiltration and the number of osteoclasts in periodontal tissues compared with the wild-type control group.³⁷ It has been reported that the level of IL-6 protein in central nervous system tissue was lower in NOX-2 knockout mice with experimental autoimmune encephalomyelitis.³⁸ In experimental atrial fibrillation mice, mRNA expressions of IL-6 and MCP-1 in atrial tissue of NOX-2 overexpressed mice showed no difference compared with wild-type control group.³⁹ These results suggest that NOX-2 may play an important role in tissue inflammation. Therefore, this study investigated the regulatory mechanism of TNF- α in cellular inflammatory response after NOX-2 gene silencing in human gingival fibroblasts. The results showed that after TNF- α stimulation of human gingival fibroblasts,




Figure 4 Changes in IL-6, MCP-1 expression level in HGFs after knockdown of NOX-2. (A) The relative mRNA expression of IL-6 was detected by RT-qPCR. (B) The relative mRNA expression of MCP-1 was detected by RT-qPCR. (C) Concentration of IL-6 in supernatant was measured by ELISA. (D) Concentration of MCP-1 in supernatant was measured by ELISA. *: $P < 0.05$; **: $P < 0.01$. si-NC: negative control siRNA group; si-NOX2: NOX-2 siRNA group.




Figure 5 Changes in ROS expression level in HGFs after knockdown of NOX-2. (A) The intracellular ROS levels with 10 ng/ml TNF- α stimulation for 48 h. Bar = 100 μ m. (B) The intracellular ROS levels that HGFs produced as measured by fluorescence microplate. **: $P < 0.01$. Abbreviations: ROS, reactive oxygen species.

the expressions of IL-6 and MCP-1 genes and proteins were significantly down-regulated in the NOX-2 gene silencing group, while the production of ROS was significantly decreased.

In conclusion, TNF- α can significantly improve the expression levels of IL-6 and MCP-1, promote the expressions of NOX-2 and ROS, and reverse the increase in the expression levels of IL-6 and MCP-1 after NOX-2 gene silencing in human gingival fibroblasts. These results suggest that TNF- α increases the expression of IL-6, MCP-1, and ROS in HGFs by upregulating the expression of NOX-2 partly.

Declaration of competing interest

The authors have no conflicts of interest relevant to this article.

Acknowledgments

This work was supported by the research funds from National Natural Science Foundation of China [grant number 82071117]; Beijing Natural Science Foundation [grant number 7212136].

References

- Bartold PM, Van Dyke TE. Periodontitis: a host-mediated disruption of microbial homeostasis. *unlearning learned concepts. Periodontology* 2013;62:203–17.
- Kwon T, Lamster IB, Levin L. Current concepts in the management of periodontitis. *Int Dent J* 2021;71:462–76.
- Sczepanik FSC, Grossi ML, Casati M, et al. Periodontitis is an inflammatory disease of oxidative stress: we should treat it that way. *Periodontology* 2020;84:45–68.
- Tóthová L, Celec P. Oxidative stress and antioxidants in the diagnosis and therapy of periodontitis. *Front Physiol* 2017;8:1055.
- Wang Y, Andrukhov O, Rausch-Fan X. Oxidative stress and antioxidant system in periodontitis. *Front Physiol* 2017;8:910.
- Tóthová L, Kamodyová N, Červenka T, et al. Salivary markers of oxidative stress in oral diseases. *Front Cell Infect Microbiol* 2015;5:73.
- Almerich-Silla JM, Montiel-Company JM, Pastor S, et al. Oxidative stress parameters in saliva and its association with periodontal disease and types of bacteria. *Dis Markers* 2015;2015:653537.
- Fredriksson MI. Effect of priming in subpopulations of peripheral neutrophils from patients with chronic periodontitis. *J Periodontol* 2012;83:1192–9.
- Aboodi GM, Goldberg MB, Glogauer M. Refractory periodontitis population characterized by a hyperactive oral neutrophil phenotype. *J Periodontol* 2011;82:726–33.
- Bedard K, Krause K. The nox family of ros-generating nadph oxidases: physiology and pathophysiology. *Physiol Rev* 2007;87:245–313.
- Buvelot H, Jaquet V, Krause KH. Mammalian nadph oxidases. 1982. In: Knaus UG, Leto TL, eds. *Methods Protoc*; 2019:17–36.
- Konior A, Schramm A, Czesnikiewicz-Guzik M, et al. Nadph oxidases in vascular pathology. *Antioxidants Redox Signal* 2014;20:2794–814.
- Lambeth JD. Nox/duox family of nicotinamide adenine dinucleotide (phosphate) oxidases. *Curr Opin Hematol* 2002;9:11–7.
- Dionigi C, Larsson L, Carcuac O, et al. Cellular expression of dna damage/repair and reactive oxygen/nitrogen species in human periodontitis and peri-implantitis lesions. *J Clin Periodontol* 2020;47:1466–75.
- Kawahara T, Ritsick D, Cheng G, et al. Point mutations in the proline-rich region of p22 are dominant inhibitors of nox1- and nox2-dependent reactive oxygen generation. *J Biol Chem* 2005;280:31859–69.
- Wang Y, Yang C. Enhanced vegf-a expression and mediated angiogenic differentiation in human gingival fibroblasts by stimulating with tnf- α in vitro. *J Dent Sci* 2022;17:876–81.
- Hajishengallis G. Immunomicrobial pathogenesis of periodontitis: keystones, pathobionts, and host response. *Trends Immunol* 2014;35:3–11.
- Frencken JE, Sharma P, Stenhouse L, et al. Global epidemiology of dental caries and severe periodontitis - a comprehensive review. *J Clin Periodontol* 2017;44:S94–105.
- Arancibia R, Oyarzún A, Silva D, et al. Tumor necrosis factor- α inhibits transforming growth factor- β -stimulated myofibroblastic differentiation and extracellular matrix production in human gingival fibroblasts. *J Periodontol* 2013;84:683–93.
- Beklen A, Al-Samadi A, Kontinen Y. Expression of cathepsin k in periodontitis and in gingival fibroblasts. *Oral Dis* 2015;21:163–9.
- Marcaccini AM, Meschiari CA, Sorgi CA, et al. Circulating interleukin-6 and high-sensitivity c-reactive protein decrease after periodontal therapy in otherwise healthy subjects. *J Periodontol* 2009;80:594–602.
- Furutani Y, Nomura H, Notake M, et al. Cloning and sequencing of the cDNA for human monocyte chemotactic and activating factor (mcaf). *Biochem Biophys Res Commun* 1989;159:249–55.
- Pype JL, Dupont LJ, Menten P, et al. Expression of monocyte chemotactic protein (mcp)-1, mcp-2, and mcp-3 by human airway smooth-muscle cells: modulation by corticosteroids and t-helper 2 cytokines. *Am J Respir Cell Mol Biol* 1999;21:528–36.
- Borch TS, Holmstrup P, Bendtzen K, et al. In vitro cytokine responses to periodontal pathogens: generalized aggressive periodontitis is associated with increased il-6 response to porphyromonas gingivalis: cytokine responses in aggressive periodontitis. *Scand J Immunol* 2010;71:440–6.
- Takashiba S, Naruishi K, Murayama Y. Perspective of cytokine regulation for periodontal treatment: fibroblast biology. *J Periodontol* 2003;74:103–10.
- Allen EM, Matthews JB, O' Halloran DJ, et al. Oxidative and inflammatory status in type 2 diabetes patients with periodontitis: periodontitis and diabetes inflammatory status. *J Clin Periodontol* 2011;38:894–901.
- Kattoor AJ, Pothineni NVK, Palagiri D, et al. Oxidative stress in atherosclerosis. *Curr Atherosclerosis Rep* 2017;19:42.
- Phull AR, Nasir B, Haq IU, et al. Oxidative stress, consequences and ros mediated cellular signaling in rheumatoid arthritis. *Chem Biol Interact* 2018;281:121–36.
- Wiegman CH, Li F, Ryffel B, et al. Oxidative stress in ozone-induced chronic lung inflammation and emphysema: a facet of chronic obstructive pulmonary disease. *Front Immunol* 2020;11:1957.
- Bañasová L, Kamodyová N, Jansáková K, et al. Salivary dna and markers of oxidative stress in patients with chronic periodontitis. *Clin Oral Invest* 2015;19:201–7.
- Buczko P, Zalewska A, Szarmach I. Saliva and oxidative stress in oral cavity and in some systemic disorders. *J Physiol Pharmacol* 2015;66:3–9.

32. Wang J, Schipper HM, Velly AM, et al. Salivary biomarkers of oxidative stress: a critical review. *Free Radic Biol Med* 2015;85: 95–104.
33. Novakovic N, Cakic S, Todorovic T, et al. Antioxidative status of saliva before and after non-surgical periodontal treatment. *Srp Arh Celok Lek* 2013;141:163–8.
34. Wang GP. Defining functional signatures of dysbiosis in periodontitis progression. *Genome Med* 2015;7:40.
35. Giannopoulou C, Krause KH, Müller F. The nadph oxidase nox2 plays a role in periodontal pathologies. *Semin Immunopathol* 2008;30:273.
36. Lin CC, Lin WN, Cho RL, et al. Tnf- α -induced cpla2 expression via nadph oxidase/reactive oxygen species-dependent nf- κ b cascade on human pulmonary alveolar epithelial cells. *Front Pharmacol* 2016;7:447.
37. Bast A, Kubis H, Holtfreter B, et al. Nadph oxidase contributes to resistance against aggregatibacter actinomycetemcomitans-induced periodontitis in mice. In: Blanke SR, ed. *Infect Immun*; 2017. 85:e00849–16.
38. Hu CF, Wu SP, Lin GJ, et al. Microglial nox2 plays a key role in the pathogenesis of experimental autoimmune encephalomyelitis. *Front Immunol* 2021;12:638381.
39. Mighiu AS, Recalde A, Zibera K, et al. Inducibility, but not stability, of atrial fibrillation is increased by nox2 over-expression in mice. *Cardiovasc Res* 2021;117:2354–64.